# An overview of chemical exposures in your environment

Lunch & Learn: EPA Finance Center 5/18/17

Laura Carlson & Jeanette Reyes\* National Center for Environmental Assessment

\*Oak Ridge institute for Science and Education Participans


#### **Conflict of Interest Statement**

- We have no conflicts of interest to disclose.
- The views expressed in this presentation are those of the author and do not necessarily reflect the views or policies of the U.S. EPA.

### Everything is a Chemical



- Pesticides/Herbicides
- Nanomaterials
- Perfluorinated Chemicals
- Plastics/Polymers
- Polycyclic Aromatic Hydrocarbons
- Flame Retardants















#### **Chemical Regulation**


- Toxic Substances Control Act (TSCA)
  - "chemicals of commerce"
  - Recently revised in 2015-2016; Lautenburg Act for Chemical Safety
- Federal Insecticide Fungicide Rodenticide Act (FIFRA)
  - Herbicides, insecticides, and pesticides
- Food Drug Administration (FDA)
  - Drugs, medical devices

#### Introduction to Toxicology

- Integrative discipline concerned with study of adverse effects of chemicals on living organisms
  - Ecotoxicology, environmental toxicology, human health
- Factors influencing toxicity
  - dosage, acute/chronic exposures, route of exposure, species, age, sex, environment
  - History: "The Dose makes the poison" -Paracelsus
- Testing methods
  - Non-human animals
  - Alternative testing methods (high throughput, modeling, etc)

#### The Chemical Problem

- Too many chemicals, not enough time/resources
  - Estimated 80,000 chemicals registered
  - Current estimates ~30,000 substances in commercial use
- Expense associated with testing
  - EPA guideline study for developmental neurotoxicity can take 1-2 years, cost \$1 million
  - Large numbers of chemicals, difficult to test with traditional methods
- Human Health Risk Assessment: function of <u>hazard identification</u>, <u>dose-response relationship</u>, <u>exposure characterization</u>, and <u>risk</u> characterization




#### **National Center for Environmental Assessment**

- Part of the Office of Research & Development
  - Split across multiple divisions: RTP, Cincinnati, Washington
- Diverse Staff
  - Biologists, chemists, ecologists, engineers, epidemiologists, toxicologists, & statisticians
- Guidance Documents / Work Products
  - Guidance documents
  - Criteria documents
  - Risk assessments
  - Risk assessment methodologies
  - Models

#### **National Center for Environmental Assessment**

- <u>human health and ecological risk assessment</u>- a robust scientific process used to determine how pollutants or other stressors may impact human health and the environment
  - interacts with other agencies, the scientific community, industry, policy-makers, and the public
  - innovative risk assessment methods and tools that help extrapolate between experimental data and real-world scenarios, improve our understanding of uncertainties, and facilitate careful evaluation of scientific evidence



#### Toxicology & Risk Assessment

- Systematic review of existing human and animal data on potential health impacts
- When sufficient data exist, development of toxicological reviews that develop recommended levels of exposure that do not come with increase risk of adverse health effects
- Look at a few case example chemicals:
  - Polychlorinated Biphenyls (PCBs)
  - Phthalates

# Polychlorinated Biphenyls (PCBs)

- 209 congeners
- Sold as commercial mixtures (1930-1977)
  - Aroclor (US); Kanechlor (Japan); Clophen (Germany), many other trade names
  - >600 million kg/yr produced in US alone
- Persistent Organic Pollutants
  - banned by TSCA 1979
  - Stockholm Convention's Dirty Dozen 2001
- Not currently in use
  - PCB contamination continues through disposal of PCB-containing products and environmental partitioning
  - Many congeners have long half lives; bioaccumulative/biomagnification


Extensive environmental contamination (ex: Great Lakes; Anniston, AL, NYC's Hudson River, etc.)

# Sources of PCB Exposure














#### Human Exposure to PCBs

- General population
  - Contaminated food (fish, meat, dairy, others)
  - Inhalation of contaminated air (indoor settings; schools)
  - ~2ng PCB/kg-d (FDA; 2003)
  - Greater Exposures-recreational fishers; native American/subsistenc
- Occupational exposures
  - Inhalation and dermal contact in workplaces where PCBs are present
- Childhood Exposures
  - · Lactational transfer through breastfeeding
  - Inhalation exposures in schools
- Difficulties with Characterizing PCB Exposures
  - Epidemiological studies use PCB serum levels, breast milk, or adipose tissue (commonly detected congeners: PCB-138, -153, and -180; PCB -28, -118, -180)
  - Exposure data consisting of only a few congeners may not accurately reflect exposures to other PCBs, which may be biologically active









#### **Emerging PCB Sources**

- Manufacturing processes result in inadvertent production of PCBs
  - Pigment production (PCB 11, 28, 52, 77, 209)
  - · Paper recycling and colored inks
- Emerging issue, high levels of lower chlorinated congeners observed in water samples; PCB-11 specifically
  - Toxicity of PCB-11 and uptake/accumulation not well understood
  - PCB-11 detected in humans
- Widespread Environmental Distribution
  - Atmospheric transport globally PCB-11
  - point sources and industrial/municipal waste water








Phthalocyanine green (pigment)

PCB-11 may be a result of direct exposure in humans; currently being evaluated by NTP

#### PCB Exposure & Health Outcomes

- PCBs have been shown to impact variety of organ systems
- The modes of action are congener-dependent
- Animal Studies evaluating PCB toxicity have observed:
  - Thyroid Effects
  - Neurological effects
  - Immunological effects
  - Reproductive effects
  - Hepatic Effects
  - Developmental Effects
  - Other organ system effects
- Currently, <u>IRIS PCB assessment (non-cancer)</u> will consider health effects listed above associated with exposure to PCB mixtures as they are found in the environment.



#### **National Exposure Research Laboratory**

- Exposure science
  - Develops tools and understanding to quantify exposure in:
    - Humans
    - Ecosystems









- Answers the questions
  - Determining risk
  - How to reduce or prevent risk
  - Are mitigation strategies successful
- Some of NERL's work is centered around
  - · Monitoring methods development
  - Exposure/dose process characterization
  - Decision support tools
  - · Predictive modeling
  - · Tools for decision making
  - Source apportionment



#### Mixtures research and cumulative assessment

- $\bullet$  Making all of those (~80,000) chemicals even more complicated...
- Chemical-by-chemical assessment
  - TSCA and REACH do not require consideration of cumulative exposures when determining human health effects
  - · Underestimate toxicity
- What happens when we look at chemicals together/jointly?
  - Additive, Synergistic, Antagonistic
- Cumulative assessment
  - WHO





#### Rapid Exposure and Dosimetry (RED) project

- RED is one of the current Chemical Safety for Sustainability (CSS) projects
  - Research projects (e.g. CSS) extend across labs/centers
- RED group
  - Human and ecological exposures for prioritization
  - develop the data, tools, and evaluation approaches required to generate rapid exposure predictions
  - models of human and ecological exposures, identification or generation of new high-throughput exposure data (e.g., chemical use or property information, consumer product use data, and consumer product and article chemical compositions, and ecological/biological monitoring data)
  - development of innovative *statistical techniques* for evaluating exposure predictions against available monitoring data.
  - develop the scientific approaches from <u>ToxCast</u> to predicted real world doses.
  - Rapid prediction allows <u>prioritization</u> based upon risk of adverse outcomes due to <u>environmental chemical exposure</u>

23


#### Understanding chemicals and quantifying exposures


- Biomonitoring data (human biological media)
  - Metabolites
  - Blood (serum and plasma), urine exhaled breath, breast milk, hair, teeth, saliva, etc.



http://www.umweitbundesamt.de/en/topics/health/commissions-working-groups/human-biomonitoring

- Physiologically Based PharmacoKinetic (PBPK)
  - PK origins in medicine
  - Time and concentration to determine dose
  - Extended to different exposures/ranges in an environ. setting
  - "PB" Based on the body compartments (mathematically modeled) instead of something entirely empirical
  - Body is arranged in a series of "compartments" set up towards a specific tissue
- High Throughput ToxicoKinetic (HTTK)
  - Large number of chemicals
  - Tying HTTK to rapid exposures






# National Health and Nutrition Examination Survey (NHANES)

- A program of studies designed to assess the <u>health</u> and <u>nutritional</u> <u>status</u> of adults and children in the United States
- The survey is unique in that it combines <u>interviews</u> and <u>p</u> examinations
- Demographics, dietary information, examination data, laboration data, questionnaire data
- Since 1999, ~5,000 people, once every 2 years, 100+ chemicals/metabolites







#### **Phthalates**


phthal-ate
/ Titalian

Social Social

Social Control

Social C

- Phthalates
  - Esters of phthalic acid used as plasticizers
  - Oily liquids are room temperature
- They are in a wide variety of goods
  - Vinyl flooring, tubing and pipes, tablecloths, pesticides, food packaging, cosmetics, skin care, some medications
- Phthalates get into the body by
  - Ingestion through food, contaminated water
  - Skin application of products on skin
  - Inhalation (from dust)
- Examples: softener for PVC pipes
- Phthalates are in everyday products
  - Metabolizes quickly but frequently exposed
  - Exposure in utero




#### **Phthalates**

| Phthalate                     | Abbreviation | Uses   | Chemical Structures  |
|-------------------------------|--------------|--|--|
| phthalic acid                 |              |  | - J  |
| di-n-butyl phthalate          | DBP          | Adhesives, caulk, cosmetics, industrial solvent                                      |  |
| diisobutyl phthalate          | DIBP         | Adhesives, caulk, cosmetics, industrial solvent                                      | Ž  |
| butyl benzyl<br>phthalate     | 88P          | Vinyl flooring, adhesives, sealants, industrial solvent                              |  |
| di(2-ethylhexyl)<br>phthalate | DEHP         | Soft plastic, including tubing, toys, home products, food containers, food packaging |  |
| diisononyl phthalate          | DINP         | Vinyt  | <b>*</b>   |
| diisodecyl phthalate          | DIDP         | plastic coating, including cook ware, pills, food packaging                          | operation of the state of the s |


#### **Phthalates**

- Some phthalates are endocrine disruptors
  - · Rat studies
  - · "phthalate syndrome"
    - · Infertility, decreased sperm count,
    - · changes in reproductive organs
- Phthalate mixture
  - Most of this group (DBP, DIBP, BBP, DINP, DEHP) associated with the "phthalate syndrome"
- How much is manufactured?
  - 90,528 tons DEHP and 2,650 tons DBP in 2012 (Lee et al., Environment International, 2014) and 235,000 tons a year (EPA 2006)
- Which have the highest toxicity of these six phthalates?
- Which can the most frequent exposures?
  - DEHP and DINP
- National Academy of Science 2008 document
  - · Cumulative risk assessment
  - investigating mechanism of action to common adverse outcome
- EPA phthalates action plan
  - Several studies have shown associations between phthalate exposures and human health
  - Under TSCA, manufacturers and processors of DPP to notify EPA at least 90 days before starting or resuming new uses of this chemical (DPP in PVC pipe)










- Differences by age and race
  - Kids (ages 6-10) have higher levels than adolescents (ages 16-17)
  - For years 2005-2008, Mexican-American children had lower levels than White non-Hispanic children and Black non-Hispanic children
  - For years 2005-2008, Black women of child-bearing age had higher concentrations than any other race



# Being An Informed Consumer

- Read product labels
- Check your products online- some NGO organizations have developed cosmetic/product database recommendations
  - Examples: Environmental Working Group's Skin Deep Database www.ewg.org
  - California Safe Cosmetics Program <a href="https://safecosmetics.cdph.ca.gov/">https://safecosmetics.cdph.ca.gov/</a>
  - Health Canada Consumer Product Safety www.hc-sc.gc.ca

#### Questions?

- Acknowledgements: NCEA co-workers
- Contact us:
  - Carlson.laura@epa.gov
  - Reves.Jeanette@epa.gov

